

# Combined EPR and DFT study of the copper(II) complexes with *N*-phosphoryl thioureas

Ruslan R. Garipov, Valery G. Shtyrin\*, Damir A. Safin, Yulia I. Zyavkina,  
Felix D. Sokolov, Alexander L. Konkin, Albert V. Aganov, Alexey V. Zakharov

Department of Chemistry, Kazan State University, Kremlevskaya 18, Kazan, 420008, Russian Federation

Received 17 April 2005; accepted 27 June 2005

Available online 28 July 2005

## Abstract

Copper(II) bis-complexes with *N*-phosphoryl thioureas,  $\text{RN}'\text{HC}(\text{S})\text{NP}(\text{O})(\text{OPr}^i)_2$  {R = phenyl (**I**), cyclohexyl (**II**)}, have been obtained and investigated in liquid and frozen toluene solutions. Equilibria between *trans*-S,N and *cis*-S,N-isomers have been revealed and described by EPR method. In contrast to more usual S,O-forms, *trans*-S,N-isomers demonstrate large superhyperfine splitting from N and P atoms simultaneously. From EPR spectra anisotropic hyperfine, superhyperfine, and *g*-tensor parameters obtained and the symmetry of the metal ion surrounding analyzed. Rhombic anisotropy of copper(II) hyperfine and axial symmetry of N and P superhyperfine splitting resolved. Structural, hydrodynamic, and magnetic characteristics of the complexes established experimentally were supported by DFT calculations. According to EPR and DFT studies, tetrahedral distortion is higher in *cis*-S,N than in *trans*-S,N forms for steric reasons and slightly higher in **I**(S,N-*trans*) isomer than in **II**(S,N-*trans*) one due to the electron-donating effect of cyclohexyl substituent. The domination of S,N- over S,O-coordination is explained by the effects of the crystal field stabilization and the formation of the intramolecular hydrogen bonds.


© 2005 Elsevier B.V. All rights reserved.

**Keywords:** EPR; DFT; Hyperfine structure; Nitrogen and phosphorus superhyperfine coupling; Copper complexes; Phosphoryl thioureas

## 1. Introduction

Copper(II) complexes with *N*-(thio)phosphoryl thioureas containing an unique pentade X–P–N–C–X in the chelate rings (where X is O or S) are very attractive to study because of possibility to relate the alterations in chelate ring atoms and the substituents at P and C atoms with the changes of dynamic and magnetic properties [1–3]. In previous X-ray and EPR studies [1–3] complexes of copper(II) with *N*-(thio)phosphoryl thioamides { $\text{RC}(\text{S})\text{NHP}(\text{X})(\text{OPr}^i)_2$ , X = S, O; R = Ph, etc.} have shown to form S,S- or S,O-coordinated molecules with six-member chelates tetrahedrally distorted from plane.

In this work we have discovered that change from  $\text{RC}(\text{S})\text{NHP}(\text{X})(\text{OPr}^i)_2$  to  $\text{RN}'\text{HC}(\text{S})\text{NHP}(\text{O})(\text{OPr}^i)_2$  {R = Ph (phenyl), Cy (cyclohexyl)} leads to an unusual coordination through S and N atoms and gives rise to the square-planar four-member metalochelates with  $\text{RN}'\text{HC}(\text{S})\text{NP}(\text{O})(\text{OPr}^i)_2$ .



An interesting feature of the compounds obtained in this work is that they exhibit resolved phosphorus superhyperfine splitting in addition to nitrogen one, what

\* Corresponding author.

E-mail address: [Valery.Shtyrin@ksu.ru](mailto:Valery.Shtyrin@ksu.ru) (V.G. Shtyrin).